

UNCLASSIFIED

AD NUMBER	
AD338814	
CLASSIFICATION CHANGES	
TO:	unclassified
FROM:	secret
LIMITATION CHANGES	
TO:	Approved for public release, distribution unlimited
FROM:	Distribution authorized to DoD and DoD contractors only; Foreign Government Information; 08 MAR 1963. Other requests shall be referred to British Embassy, 3100 Massachusetts Avenue, NW, Washington, DC 20008.
AUTHORITY	
DSTL, WO 195/15601, 20 May 2008; DSTL, WO 195/15601, 20 May 2008	

THIS PAGE IS UNCLASSIFIED

**SECRET**

**AD** **338 814L**

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



**SECRET**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

NOTICE:

THIS DOCUMENT CONTAINS INFORMATION  
AFFECTING THE NATIONAL DEFENSE OF  
THE UNITED STATES WITHIN THE MEAN-  
ING OF THE ESPIONAGE LAWS, TITLE 18,  
U.S.C., SECTIONS 793 and 794. THE  
TRANSMISSION OR THE REVELATION OF  
ITS CONTENTS IN ANY MANNER TO AN  
UNAUTHORIZED PERSON IS PROHIBITED  
BY LAW.

SECRET-DISCREET

P.T.P. 853

P.T.P. 853

Copy No. 65

PORTON TECHNICAL PAPER No 853

CATALOGED BY DDC  
AS AD No. 338814

AN ESTIMATE OF THE HUMAN INHALATION

L(Ct)50 FOR GB [U]

1. THE INFORMATION CONTAINED IN THIS DOCUMENT IS UNCLASSIFIED EXCEPT WHERE SHOWN OTHERWISE BY THE MARKING IN ANY OTHER MANNER. IT IS THE POLICY OF THE GOVERNMENT OF THE UNITED KINGDOM TO RELEASE AS MUCH OF THIS INFORMATION AS POSSIBLE.

BY

2. THE INFORMATION SHOULD BE SAFEGUARDED UNDER RULES DESIGNED TO PROTECT THE SAME STATUS AS THE SECURITY AS THAT IS ENJOYED BY HER MAJESTY'S GOVERNMENT IN THE UNITED KINGDOM.

3. THE INFORMATION CONTAINED IN THIS DOCUMENT SHOULD NOT BE DISCLOSED OUTSIDE GOVERNMENT DEPARTMENTS WITHOUT THE PRIOR PERMISSION OF THE MINISTRY OF DEFENCE.

4. THE RECIPIENT IS WARNED THAT INFORMATION CONTAINED IN THIS DOCUMENT MAY BE SUBJECT TO PROMOTE THE INTERESTS OF THE UNITED KINGDOM.

R.J. SHEPHARD

DDC  
RECEIVED  
AUG 5 1963  
TISIA D

CHEMICAL DEFENCE EXPERIMENTAL ESTABLISHMENT

338 814L

Porton, Wilts.

EXCLUDED FROM AUTOMATIC  
RECLASSIFICATION: DOD DIR 5200.10  
Does Not Apply

SECRET-DISCREET

**SECRET**

THIS DOCUMENT IS THE PROPERTY OF H.B.M. GOVERNMENT,  
and is issued for the information only of those  
officials who are concerned with its contents.

The official in possession of the document will be  
responsible for its safe custody and when not in use it  
is to be kept under lock and key.

SECRET-DISCREET

PORTON TECHNICAL PAPER NO. 853

COPY NO. 65

DATE: 8th March, 1963.

AN ESTIMATE OF THE HUMAN INHALATION L(Ct)50  
FOR GB

By

R.J. Shephard

SUMMARY

1. Methods of estimating the inhalation L(Ct)50 are outlined.
2. Reasons are suggested for basing calculations on the ratio ChE50: ID50; this gives an estimate of the average L(Ct)50 that is probably valid to within  $\pm 33\%$  for brief exposures, with wider limits for longer exposures.
3. The L(Ct)50 varies with the metabolic state of those attacked. Under resting conditions, the average figure is probably at least 135 ng.min/m<sup>3</sup>; with activity the average may be less than 50 ng.min/m<sup>3</sup>.
4. These figures represent the best possible estimate based on data available from both U.S. and the U.K.

This document contains information affecting the National Defense of the United States within the meaning of the Espionage Laws, Title 18, U. S. C., Section 793 and 794. Its transmission or the revelation of its contents in any manner to an unauthorized person is prohibited by law.

RJS/PMT. C. Lovatt Evans,  
Head, Physiology Section.  
(Sgd.) W.S.S. Ladell,  
Assistant Director(Medical).

SECRET-DISCREET

SECRET-DISCREET

PORTON TECHNICAL PAPER NO.853
COPY NO. 65
DATE. 8th March, 1963.

AN ESTIMATE OF THE HUMAN INHALATION L(Ct)50  
FOR GB

By

R.J. Shephard

1. HISTORICAL. The human inhalation L(Ct)50 has usually been calculated from an assumed ventilation (10 l/min in the resting state) and an LD50 ( $\mu\text{g GB/kg}$ ) obtained by one of several methods of extrapolation. Complete retention of inhaled GB has usually been assumed.

- (a) The U.K. approach. Ainsworth, Davies and McKee (1) gave single-breath inhalations of GB ( $\frac{1}{2}$  - 3  $\mu\text{g/kg}$ ) to 36 subjects. The dose of GB was plotted against the logarithm of the percentage of uninhibited red cell enzyme, and by considerable extrapolation of the graph, the dose for 90% inhibition was estimated at 11 - 29  $\mu\text{g/kg}$ , with a most probable value of 16  $\mu\text{g/kg}$ . The assumptions of the method were
- (i) that the dose of GB causing 90% inhibition of red cell cholinesterase was the LD50, and
  - (ii) that the dose of GB was linearly related to the logarithm of the percentage of enzyme remaining after administration of GB. The last assumption was checked by giving GB to rabbits (inhalation and i.v.) and guinea pigs (s.c.) over a wider dose range. No gross departure from this relationship was apparent, but the number of observations was not adequate to study the fine details of the curve.

SECRET-DISCREET

SECRET-DISCREET

Assuming a ventilation of 10 l/min, with 100% retention of GB, the resting L(Ct)50 was placed in the range 77 - 203 mg.min/m<sup>3</sup>, with a most probable value of 112 mg.min/m<sup>3</sup>. It is now known that assumption (1) is not universally true. Many species of animals remain active when the red cell cholinesterase is 90% inhibited, and recent U.S. experiments suggest the same is true of man. If a figure of 95% inhibition is taken, the estimated L(Ct)50 would be 224 mg.min/m<sup>3</sup>.

- (b) The U.S. approach Silver (2) considered that during short exposures to GB vapour, the inhalation LD50 for a number of laboratory animals approached the intravenous LD50 of about 15 µg/kg. It was assumed that the same would be true of man, and again accepting a respiratory minute volume of 10 l/min, with 100% retention of agent, the L(Ct)50 would be 105 mg.min/m<sup>3</sup> for a 70 kg man.

2. PRESENT POSITION. The basic method of estimation has altered little, but much more information is now available on the response of man to moderate doses of GB by inhalation, on the respiratory minute volume during exposure and simulated exposure to "nerve gas", and on the retention of inhaled vapours.

- (a) Extrapolation to LD50. The estimation of the LD50 remains the weakest link in the chain of evidence. To the original two methods of extrapolation has been added a third, based on the ratio of the ChE50 to the LD50; this seems the procedure of choice, for the reasons discussed below.

- (1) Cholinesterase inhibition at LD50. Whether log-probit (3) or semi-logarithmic plot of the data is used, the extrapolation is critically dependent on the slope, which in man can only be determined over a very small part of the dose range of interest; further, there is no certain evidence that 90% is the desired inhibition. From the data of Callaway, Davies and Rutland (4), in many species the LD50 dose corresponds to >95% inhibition of whole blood and red cell cholinesterase:

SECRET-DISCREET



SECRET-DISCREET

Species	LD50 ( $\mu\text{g/kg}$ , s.c.)	Corresponding cholinesterase inhibition	
		Whole blood	Red cell
Rabbit	35	97	> 99.9
Monkey	25	97	
Guinea Pig	38	98	> 99.9
Rat	109	91	
Pigeon	52	88	

(11) Inter-species comparison of LD50: There is a wide range of species variation in inhalation toxicity of GB, and the supposition that inhalation toxicity approaches intravenous toxicity (2) is not fully borne out by experimental data. The inhalation toxicity apparently increases with a decrease in the duration of exposure (5 - 8); however, even with short exposures (9 - 10), the intravenous LD50 is exceeded by a factor of 1.7 - 2.0

Species	Inhalation LD50 $\mu\text{g/kg}$	Intravenous LD50 $\mu\text{g/kg}$
Rat (4)	165 - 209	80
Guinea Pig (4)	45 - 73	32
Guinea Pig (9)	42 - 79	32 - 36
Rabbit (4)	25 - 52	19
Monkey (4)	46 - 48	-
Monkey (9)	15 - 38	18 - 23
Sheep (4)	58 - 76	15 - 20
Dog (9)	28 - 52	18 - 28

SECRET - DISCREET

SECRET-DISCREET

- (iii) Ratio of ChE50 to LD50. The ChE50: LD50 ratio has been thought rather constant in different species. U.K. data (4) show a range of 3.2 - 6.3, with a mean value of about 4.0:

Species	ChE50 : LD50 ratio	
	Whole blood	Red cell
Rabbit	1: 6.0	1 : 6.3
Monkey	1: 6.3	
Guinea Pig	1: 3.3	1 : 3.4
Rat	1: 3.2	
Pigeon	1: 4.0	

U.S. experiments show a mean value 3.9 (11). The use of the ChE50 : LD50 ratio as a basis of extrapolation seems preferable to methods (i) and (ii). The quantities involved (ChE50 and LD50 in animals, ChE50 in man) can all be measured accurately, and no assumptions are made about the shape of the cholinesterase inhibition curve in the range 90 - 95% inhibition, which is difficult to check experimentally. The influence of route of administration on the ChE50/LD50 ratio is less than might be anticipated, changes in LD50 being matched at least in part of changes in ChE50. Thus, the following ratios have been reported for the rabbit:

Route of administration	LD50 (µg/kg)	Whole blood ChE at LD50	ChE 50 : LD50 ratio
Subcutaneous (4)	35	97	1 : 6.0
Intravenous (9, 3)	19	94	1 : 3.4
Inhalation (9, 1)	25 - 52	97 - 100	1 : 5.2 to 1 : 10.8

SECRET-DISCREET

SECRET-DISCREET

In the case of the inhalation experiments, the ratio increased as the duration of exposure was lengthened. With large doses of GB, the cholinesterase inhibition at the ID50 also varies with exposure time. Thus some figures of Cullumbine, Callaway, Ainsworth and Lynch (12) show that in the sheep the ID50 with 2 min exposure is 76  $\mu\text{g/kg}$ , with 90% cholinesterase inhibition, and with 10 sec exposure the ID50 is 58  $\mu\text{g/kg}$ , with 95.9% inhibition of whole blood cholinesterase. This difference is sufficient to preclude effective use of prediction method (1). On the other hand, the ChE50 dose seems relatively independent of the duration of exposure (13, 14).

- (b) The inhalation red cell ChE50 in man. Whether based on very brief exposures (single-breath technique, 2 sec or less) or 15 min chamber exposures, the ChE50 can be placed fairly certainly between 4.0 and 5.0  $\mu\text{g/kg}$ :

Method of inhalation	Method of extrapolation	Number of subjects	Estimated red cell ChE50 ( $\mu\text{g/kg}$ )
Single breath (1)	Semi-log plot	36	4.7
" " (14)	Linear regression	73	5.0*
" "	log/probit plot(3)	29 paired obs.	4.0*
		131 obs.	4.4*
Chamber			
15 min seated }	log/probit plot(3)	92 }	5.0
15 min marching }		44 }	

In the chamber exposures, the inhaled dose has been back-calculated (14) from the Ct and the ventilation measured in parallel experiments, using 96% retention for the resting experiments (15) and 90% (10, 16) for the exercising men.

\*In these calculations, 67% recovery of nominal dose assumed (14).

SECRET-DISCREET

SECRET-DISCREET

- (c) The inhalation LD50. If man can be assumed to behave more like a monkey than some of the small species, the ChE50 : LD50 ratio of 1 : 4.0 may be rather low. A better estimate might be 1 : 5 for short exposures ( $\leq \frac{1}{2}$  min) and 1 : 11 for longer exposures ( $> 2$  min). On this basis, the inhalation LD50 would be:

Duration	LD50	Total dose for 70 kg man
$\leq \frac{1}{2}$ min	20 - 25 $\mu\text{g/kg}$	1400 - 1750 $\mu\text{g}$
$> 2$ min	44 - 55	3080 - 3950

The influence of duration of exposure on the LD50 remains somewhat uncertain. It is clear from the figures in section (b), above, that at doses likely to produce  $< 50\%$  inhibition of circulating cholinesterase, the effectiveness of a given inhaled dose is similar during both "single-breath" and 15 min chamber exposures. However, with larger doses, the probability of extravascular spread will rise with increase of concentration of the agent in the blood stream, and for this reason the toxicity of a given large dose may be greater when it is inhaled rapidly. The total dose corresponding to the LD50 will rise with body weight; however, this will be offset to some extent by corresponding differences of respiratory minute volume, and over a small range of body weights differences in  $L(Ct)_{50}$  from this factor can probably be neglected.

SECRET-DISCREET

SECRET-DISCREET

- (d) Percentage retention of inhaled GB. Direct measurement has shown 96% retention of GB during nasal breathing (15). Calculation suggests 88 - 90% for oral breathing (16); U.S. figures are a little lower (83 - 88%, ref. 9). Thus we may assume nasal breathing (96% retention) at rest, and a combination of oral and nasal breathing in exercise (say 90% retention).
- (e) Respiratory minute volume. With exposures <15 sec, the breath may be held. With longer exposures ventilation will gradually come to equal the metabolic requirement of the activity undertaken. Typical figures for the period  $\frac{1}{2}$  - 20 min are:

Activity	Range of minute volume l/min	Average l/min
Resting, seated	10 - 14	12
" standing	12 - 20	16
Marching 2.5 m.p.h.	18 - 60	29
Running (max. effort)	60 - 100	80

- (f) The inhalation L(Ct)50. Combining the information in (e), (d) and (e), the estimated L(Ct)50 for a 70 kg man may be tabulated as follows:

Activity	Exposure $\frac{1}{2}$ min		Exposure 2 min	
	Range mg.min/m <sup>3</sup>	Average	Range	Average
Resting, seated	104 - 182	135	230 - 410	301
" standing	73 - 152	102	161 - 343	230
Marching 2.5 m.p.h.	26 - 108	60	57 - 244	135
Running (max. effort)	16 - 32	22	34 - 73	49

SECRET-DISCREET

SECRET-DISCREET

### 3. CONCLUSIONS

Although it is possible for planning purposes to tabulate the anticipated L(Ct)50 at various levels of metabolism, as above, there remain important uncertainties in the estimate. The ratio CHE50 : LD50 shows a twofold variation between species, and it is by no means certain which species man resembles. It is also quite probable that whereas the ratio found for brief inhalation exposures corresponds with the intravenous ratio, the ratio applicable to longer inhalation exposures is greater than the intravenous ratio. The average figures quoted above could thus still be in error by about a third in the case of brief exposures, and by a larger margin for longer exposures.

(Sgd.) C. Lovatt Evans,  
Head, Physiology Section.

(Sgd.) W.S.S. Ledell,  
Assistant Director(Medical).

RJS/PNF.

SECRET-DISCREET

SECRET-DISCREET

REFERENCES

1. Ainsworth, M., Davies, D.R. and McKee, W.H.E. (1951) Ptn./TA.2620/34.61/51.
2. Silver, S.D. (1953), M.L.S.R. 23
3. 15th Tripartite Meeting, Medical Aspects. U.K. Discussion Paper TC/15/11
4. Callaway, S., Davies, D.R. and Rutland, J.P. (1954), P.T.P. 425.
5. McDonald, F. (1945), Addendum to P.R. 2693.
6. Barrett, H.M. and McPhail, M.K. (1949), S.T.M. 132
7. McPhail, M.K. and Barrett, H.M. (1950), D.R.B. Symposium.
8. Krakow, E.H. (1948), M.D.R. 142
9. Cullumbine, H. (1956), Porton Report.
10. Oberst, F.W. (1960, British Occupational Hygiene Society Symposium Oxford.
11. Marsulli, F.N. and Callahan, J.F. (1954), M.L.R.R. 249
12. Cullumbine, H., Callaway, S., Ainsworth, M. and Lynch, R. (1955), P.T.P. 495.
13. Whitchoer, H.W. (1959), P.T.P. 695.
14. Shephard, R.J. (1959), P.T.P. 702.
15. Ainsworth, M. and Truckle, T.W.N. (1952), P.T.P. 244.
16. Shephard, R.J. and Ainsworth, M. (1961), P.T.P. 773.

SECRET-DISCREET

SECRET-DISCREET

DISTRIBUTION

to Porton Technical Paper No. 853

WAR OFFICE

1 Chief Scientist  
2 D.C.S.(A)  
3 -6 D.C.D.R.D.  
7 D.P.B.R.  
8 -9 E.P.4

R. & D. ESTABLISHMENTS

84-109 C.D.E.E.  
10 A.O.R.E. (F)

ADVISORY BODIES

Chemical Defence Advisory Board

11 Prof. R.B. Fisher  
12 Prof. N.K. Adam  
13 Dr. J.M. Barnes  
14 Dr. O.G.  
15 Prof. D.H. Hey  
16 Prof. B.J. Mason  
17 Prof. H.N. Rydon  
18 Mr. A.L. Stuchbery  
19 Prof. D.D. Woods

Biology Committee

20 Prof. E. Boyland  
21 Lord Evans  
22 Prof. W.T. Irvine  
23 Prof. Sir Aubrey Lewis  
24 Dr. R.D. Keynes  
25 Sir Charles Lovatt Evans  
26 Prof. W.D.M. Paton  
27 Sir Rudolph Peters  
28 Dr. J.R. Squire  
29-38 Secretariat S.A.C.

MINISTRY OF AVIATION

39 Pats.1  
40-41 T.I.L.

U.K. HIGH COMMISSION, OTTAWA

42 Senior Army Liaison Officer

BRITISH DEFENCE STAFFS, WASHINGTON

43-49 R. Holmes, Esq., D.R. Staff

OVERSEAS (through T.I.L.)

Australia

50-52 Defence Standard Laboratories  
53 Senior Representative, Dept. of Supply  
54 Army Staff (U.K.)  
55 R.A.A.F. (Tech. Section)

Canada

56-57 Chairman, Defence Research Board  
58-59 Defence Research Chemical Laboratories,  
Ottawa.  
60 Suffield Experimental Station

U.S.A.

61-73 Reading Panel  
74-80 U.S. Chem. Corps Liaison Officer,  
Porton.

SECRET-DISCREET





*Information Centre  
Knowledge Services  
[dstl] Porton Down,  
Salisbury  
Wiltshire  
SP4 0JQ  
22060-6218  
Tel: 01980-613753  
Fax 01980-613970*

Defense Technical Information Center (DTIC)  
8725 John J. Kingman Road, Suit 0944  
Fort Belvoir, VA 22060-6218  
U.S.A.

AD#: AD338814

Date of Search: 20 May 2008

Record Summary: WO 195/15601

Title: Estimate of the Human Inhalation L(CT)50 for GB  
Availability Open Document, Open Description, Normal Closure before FOI Act: 30 years  
Former reference (Department) SAC-676  
Held by The National Archives, Kew

This document is now available at the National Archives, Kew, Surrey, United Kingdom.

DTIC has checked the National Archives Catalogue website (<http://www.nationalarchives.gov.uk>) and found the document is available and releasable to the public.

Access to UK public records is governed by statute, namely the Public Records Act, 1958, and the Public Records Act, 1967.

The document has been released under the 30 year rule.

(The vast majority of records selected for permanent preservation are made available to the public when they are 30 years old. This is commonly referred to as the 30 year rule and was established by the Public Records Act of 1967).

This document may be treated as **UNLIMITED**.